Author Index

Abramson, D.
Measurement of ochratoxin A in barley extracts by liquid chromatography–mass spectrometry 315
Allain, P., see Cailleux, A. 280
Aue, W. A., see Siu, K. W. M. 433
Beger, J., see Voelkel, A. 373
Bergman, D. A.
—— and Winzor, D. J.
Allowance for "analyte valence" in the retention model of ion-exchange chromatography. Adenosine 5′-phosphates on DEAE cellulose 67
Berlizov, Yu. S., see Buryan, P. 89
Berman, S. S., see Siu, K. W. M. 433
Blakemore, W. M.
——, Rushing, L. G., Thompson, Jr., H. C., Freeman, J. P., Levine, R. A. and Nony, C. R.
Characterization, purification, and analysis of Solvent Yellow 33 and Solvent Green 3 dyes 219
Bond, A. D., see Melucci, C. K. 321
Borén, H., see Wigilius, B. 169
Bossche, W. Van den, see De Schutter, J. A. 303
Burden, R. S.
——, Deas, A. H. B. and Clark, T.
Separation of enantiomers of fungicidal and plant growth regulatory triazole alcohols using chiral derivatisation and capillary gas chromatography 273
Buryan, P.
——, Macák, J., Třiska, J., Vodička, L., Berlizov, Yu. S., Dmitrikov, V. P. and Nabivach, V. M.
Kováts retention indices of alkylquinolines on capillary columns 89
Cabras, P.
——, Meloni, M. and Pirisi, F. M.
Separation of Benalaxyl and its metabolites by high-performance liquid chromatography 334
Cailleux, A.
——, Turcant, A., Allain, P., Toussaint, D., Gaste, J. and Roux, A.
Gas chromatographic analysis of volatile compounds in water and biological samples with an automatic injector 280
Carlberg, G. E., see Wigilius, B. 169
Chan, K. C.
—— and Yeung, E. S.
Comments on enhanced peak response due to solvent interaction 465
Chaudhury, M. K., see Van Oss, C. J. 53
Chung, G. N., see Reymond, D. 97
Clark, T., see Burden, R. S. 273
Čoupek, J., see Vláčil, F. 119
Deas, A. H. B., see Burden, R. S. 273
Deeble, S., see Sidhu, A. S. 233
De Moerloose, P., see De Schutter, J. A. 303
De Schutter, J. A.
——, Van den Bossche, W. and De Moerloose, P.
Stability-indicating analysis of tetryzoline hydrochloride in pharmaceutical formulations by reversed-phase ion-pair liquid chromatography 303
Dewaele, C., see Verzele, M. 111
Dmitrikov, V. P., see Buryan, P. 89
Drevin, I., see Johansson, B.-L. 448
Duquet, D., see Verzele, M. 111
Ebert, K., see Voelkel, A. 373
Engelhardt, R., see Gabius, H.-J. 452
Folestad, S.
——, Josefsson, B. and Larsson, M.
Performance and preparation of immobilized polysiloxane stationary phases in 5–55 μm I.D. open-tubular fused-silica columns for liquid chromatography 347
Freeman, J. P., see Blakemore, W. M. 219
Gabius, H.-J.
—— and Engelhardt, R.
Interaction of mannose-binding proteins with different types of immobilized affinity ligands 452
Gaste, J., see Cailleux, A. 280
Gill, R.
—— and Wanogho, S. O.
High-performance liquid chromatographic system for the separation of tricyclic antidepressant and related drugs using ODS-Hypersil 461
Gont, L. K.
—— and Neuendorf, S. K.
Enantiomeric separation of N-carbamyltryptophan by thin-layer chromatography on a chiral stationary phase 343
Good, R. J., see Van Oss, C. J. 53
Goto, M., see Ito, Y. 296
Grimvall, A., see Wigilius, B. 169
Grob, K.
—— and Schilling, B.
Uncoated capillary column inlets (retention gaps) in gas chromatography (Review) 3
Haddad, P. R., see Ivey, J. P. 309
Hadziyev, D., see Nguyen, T. T. 257

Author Index

Abramson, D.
Measurement of ochratoxin A in barley extracts by liquid chromatography–mass spectrometry 315
Allain, P., see Cailleux, A. 280
Aue, W. A., see Siu, K. W. M. 433
Beger, J., see Voelkel, A. 373
Bergman, D. A.
—— and Winzor, D. J.
Allowance for "analyte valence" in the retention model of ion-exchange chromatography. Adenosine 5′-phosphates on DEAE cellulose 67
Berlizov, Yu. S., see Buryan, P. 89
Berman, S. S., see Siu, K. W. M. 433
Blakemore, W. M.
——, Rushing, L. G., Thompson, Jr., H. C., Freeman, J. P., Levine, R. A. and Nony, C. R.
Characterization, purification, and analysis of Solvent Yellow 33 and Solvent Green 3 dyes 219
Bond, A. D., see Melucci, C. K. 321
Borén, H., see Wigilius, B. 169
Bossche, W. Van den, see De Schutter, J. A. 303
Burden, R. S.
——, Deas, A. H. B. and Clark, T.
Separation of enantiomers of fungicidal and plant growth regulatory triazole alcohols using chiral derivatisation and capillary gas chromatography 273
Buryan, P.
——, Macák, J., Třiska, J., Vodička, L., Berlizov, Yu. S., Dmitrikov, V. P. and Nabivach, V. M.
Kováts retention indices of alkylquinolines on capillary columns 89
Cabras, P.
——, Meloni, M. and Pirisi, F. M.
Separation of Benalaxyl and its metabolites by high-performance liquid chromatography 334
Cailleux, A.
——, Turcant, A., Allain, P., Toussaint, D., Gaste, J. and Roux, A.
Gas chromatographic analysis of volatile compounds in water and biological samples with an automatic injector 280
Carlberg, G. E., see Wigilius, B. 169
Chan, K. C.
—— and Yeung, E. S.
Comments on enhanced peak response due to solvent interaction 465
Chaudhury, M. K., see Van Oss, C. J. 53
Chung, G. N., see Reymond, D. 97
Clark, T., see Burden, R. S. 273
Čoupek, J., see Vláčil, F. 119
Deas, A. H. B., see Burden, R. S. 273
Deeble, S., see Sidhu, A. S. 233
De Moerloose, P., see De Schutter, J. A. 303
De Schutter, J. A.
——, Van den Bossche, W. and De Moerloose, P.
Stability-indicating analysis of tetryzoline hydrochloride in pharmaceutical formulations by reversed-phase ion-pair liquid chromatography 303
Dewaele, C., see Verzele, M. 111
Dmitrikov, V. P., see Buryan, P. 89
Drevin, I., see Johansson, B.-L. 448
Duquet, D., see Verzele, M. 111
Ebert, K., see Voelkel, A. 373
Engelhardt, R., see Gabius, H.-J. 452
Folestad, S.
——, Josefsson, B. and Larsson, M.
Performance and preparation of immobilized polysiloxane stationary phases in 5–55 μm I.D. open-tubular fused-silica columns for liquid chromatography 347
Freeman, J. P., see Blakemore, W. M. 219
Gabius, H.-J.
—— and Engelhardt, R.
Interaction of mannose-binding proteins with different types of immobilized affinity ligands 452
Gaste, J., see Cailleux, A. 280
Gill, R.
—— and Wanogho, S. O.
High-performance liquid chromatographic system for the separation of tricyclic antidepressant and related drugs using ODS-Hypersil 461
Gont, L. K.
—— and Neuendorf, S. K.
Enantiomeric separation of N-carbamyltryptophan by thin-layer chromatography on a chiral stationary phase 343
Good, R. J., see Van Oss, C. J. 53
Goto, M., see Ito, Y. 296
Grimvall, A., see Wigilius, B. 169
Grob, K.
—— and Schilling, B.
Uncoated capillary column inlets (retention gaps) in gas chromatography (Review) 3
Haddad, P. R., see Ivey, J. P. 309
Hadziyev, D., see Nguyen, T. T. 257

Harada, K.
——, Nishimura, O. and Mihara, S.
Rapid analysis of coffee flavour by gas chromatography using a pyrolyzer 457
Hashimoto, T., see Kato, Y. 395
Hormazabal, V., see Rogstad, A. 328
Hsu, T.-B.
——, Shah, P. A. and Rogers, L. B.
Synthesis and characterization of chiral stationary phases from amino acids and small peptides for liquid chromatography fractionations of a racemic alcohol 145
Ikeda, A., see Iwahashi, H. 199
Ishibashi, M.
——, Watanabe, K., Yamashita, K., Miyazaki, H. and Krolik, S.
Diethylhydrogensilyl-cyclic diethylsilylene derivatives in gas chromatography–mass spectrometry of prostanoids. III. F_α-prostaglandins and thromboxane B_2 derivatives 183
Ishii, D., see Ito, Y. 296
Ito, Y.
—— and Lee, Y. W.
Analytical high-speed counter-current chromatography with a coil planet centrifuge 290
——, Takeuchi, T., Ishii, D., Goto, M. and Mizuno, T.
Direct coupling of micro high-performance liquid chromatography with fast atom bombardment mass spectrometry. III. Application to oligosaccharides 296
Ivey, J. P.
—— and Haddad, P. R.
Determination of dimethylsulphoxide using ion-exclusion chromatography with ultraviolet absorption detection 309
Iwahashi, H.
——, Negoro, Y., Ikeda, A. and Kido, R.
Detection of some retinoid radicals using high-performance liquid chromatography with electron spin resonance spectroscopy or electrochemical detection 199
James, L. B.
Amino acid analysis: a replacement solvent for flushing ninhydrin reagent 269
Janák, J., see Soják, L. 79
Johansson, B.-L.
—— and Drevin, I.
Quantitative determination of the ligand in Phenyl-Sepharose FF with proton nuclear magnetic resonance and derivative ultraviolet spectroscopy 448
Johnson, R. N., see Melucci, C. K. 321
Jokl, V.
——, Polášek, M. and Pospíchalová, J.
Isotachophoretic determination of the ionic mobilities and ionization constants of weak monoacidic bases by a simple computer-aided slope–intercept method 427
Josefsson, B., see Folestad, S. 347
Kato, Y.
——, Kitamura, T., Nakamura, K., Mitsui, A., Yamasaki, Y. and Hashimoto, T.
High-performance liquid chromatography of membrane proteins 395
Kawai, S.
——, Sato, Y., Takagi, S.-I. and Nomoto, K.
Separation and determination of mugineic acid and its analogues by high-performance liquid chromatography 325
Kennedy, J. M., see Sidhu, A. S. 233
Kido, R., see Iwahashi, H. 199
Kitamura, T., see Kato, Y. 395
Kokkonen, P.
——, Ojaniemi, E. and Pöykiö, R.
Analysis of coke oven gas by gas chromatography 435
Krolik, S., see Ishibashi, M. 183
Lafuente, M. T., see Tadeo, J. L. 338
Larsson, M., see Folestad, S. 347
Lee, Y. W., see Ito, Y. 290
Levine, R. A., see Blakemore, W. M. 219
Lundgren, B. V., see Wigilius, B. 169
Lyman, G. W., see Melucci, C. K. 321
Macák, J., see Buryan, P. 89
Maitani, T.
——, Uchiyama, S. and Saito, Y.
Hydride generation–flame atomic-absorption spectrometry as an arsenic detector for high-performance liquid chromatography 161
Martire, D. E., see Schantz, M. M. 35
Mayer, J. M., see Reymond, D. 97
Meier, B., see Ziegler, S. J. 419
Meloni, M., see Cabras, P. 334
Melucci, C. K.
——, Lyman, G. W., Bond, A. D. and Johnson, R. N.
Determination of *S*-carboxymethylcysteine in syrup formulations by high-performance liquid chromatography 321
Meyer, H.-P., see Neumann, H. 442
Mihara, S., see Harada, K. 457
Mitsui, A., see Kato, Y. 395
Miyazaki, H., see Ishibashi, M. 183
Mizuno, T., see Ito, Y. 296
Moerloose, P., De, see De Schutter, J. A. 303
Nabivach, V. M., see Buryan, P. 89
Nakamura, K., see Kato, Y. 395

Nawrocki, J.
Influence of silica surface heterogeneity on the retention of hydrocarbons in gas chromatography 266
Negoro, Y., see Iwahashi, H. 199
Neuendorf, S. K., see Gont, L. K. 343
Neumann, H.
—— and Meyer, H.-P.
Application of a gas chromatographic capillary-to-capillary column-switching system to the analysis of complex illicit drug samples 442
Nguyen, T. T.
——, Palcic, M. M. and Hadziyev, D.
High-performance liquid chromatographic assay of potato tuber 3′-nucleotidase 257
Nishimura, O., see Harada, K. 457
Nomoto, K., see Kawai, S. 325
Nony, C. R., see Blakemore, W. M. 219
Ojaniemi, E., see Kokkonen, P. 435
Oss, C. J. van, see Van Oss, C. J. 53
Palcic, M. M., see Nguyen, T. T. 257
Pirisi, F. M., see Meloni, M. 334
Pöykiö, R., see Kokkonen, P. 435
Polášek, M., see Jokl, V. 427
Pospíchalová, J., see Jokl, V. 427
Pussayanawin, V.
—— and Wetzel, D. L.
High-performance liquid chromatographic determination of ferulic acid in wheat milling fractions as a measure of bran contamination 243
Reymond, D.
——, Chung, G. N., Mayer, J. M. and Testa, B.
Lipophilicity measurement of nicotinates by reversed-phase high-performance liquid chromatography. Differences in retention behaviour, but similarities of log k_w values, in methanol–water and acetonitrile–water eluents 97
Rogers, L. B., see Hsu, T.-B. 145
Rogstad, A.
——, Hormazabal, V. and Yndestad, M.
Extraction of praziquantel from fish tissue and its determination by high-performance liquid chromatography 328
Roux, A., see Cailleux, A. 280
Roy, S.
High-performance liquid chromatographic analysis of chloropigments (Review) 19
Ruman, J., see Soják, L. 79
Rushing, L. G., see Blakemore, W. M. 219
Sävenhed, R., see Wigilius, B. 169
Saito, Y., see Maitani, T. 161
Sato, Y., see Kawai, S. 325
Schantz, M. M.
—— and Martire, D. E.
Determination of hydrocarbon–water partition coefficients from chromatographic data and based on solution thermodynamics and theory 35
Schilling, B., see Grob, K. 3
Schutter, J. A. De, see De Schutter, J. A. 303
Shah, P. A., see Hsu, T.-B. 145
Shoji, T., see Takatsuki, K. 207
Sidhu, A. S.
——, Kennedy, J. M. and Deeble, S.
General method for the analysis of pharmaceutical dosage forms by high-performance liquid chromatography 233
Siu, K. W. M.
——, Berman, S. S. and Aue, W. A.
Response correlation among d.c. and low-frequency a.c. regimes in electron-capture detection 433
Soják, L.
——, Ruman, J. and Janák, J.
Characterization of monoalkylcyclopentadiens by retention–structure correlation in capillary gas chromatography 79
Sticher, O., see Ziegler, S. J. 419
Szymanowski, J., see Voelkel, A. 373
Tadeo, J. L.
—— and Laufente, M. T.
Determination of fenpropimorph in citrus fruit by reversed-phase high-performance liquid chromatography 338
Takagi, S.-I., See Kawai, S. 325
Takatsuki, K.
——, Ushizawa, I. and Shoji, T.
Gas chromatographic–mass spectrometric determination of macrolide antibiotics in beef and pork using single ion monitoring 207
Takeuchi, T., see Ito, Y. 296
Testa, B., see Reymond, D. 97
Thompson, Jr., H. C., see Blakemore, W. M. 219
Toussaint, D., see Cailleux, A. 280
Tříska, J., see Buryan, P. 89
Turcant, A., see Cailleux, A. 280
Uchiyama, S., see Maitani, T. 161
Ushizawa, I., see Takatsuki, K. 207
Van den Bossche, W., see De Schutter, J. A. 303
Van Oss, C. J.
——, Good, R. J. and Chaudhury, M. K.
Mechanism of DNA (Southern) and protein (Western) blotting on cellulose nitrate and other membranes 53
Verzele, M.
——, Dewaele, C. and Duquet, D.
Observations and ideas on slurry packing of liquid chromatography columns 111
——, see Yang, Y.-B. 383

Vinš, I., see Vláčil, F. 119, 133
Vláčil, F.
—— and Vinš, I.
Modified hydroxyethyl methacrylate copolymers as sorbents for ion chromatography. II. Influence of the functional group on sorbent selectivity 133
——, Vinš, I. and Čoupek, J.
Modified hydroxyethyl methacrylate copolymers as sorbents for ion chromatography. I. Synthesis and properties of sorbents 119
Vodička, L., see Buryan, P. 89
Voelkel, A.
——, Szymanowski, J., Beger, J. and Ebert, K.
Polarity of aminoether alcohols and their ethers measured by reversed-phase gas chromatography 373
Wanogho, S. O., see Gill, R. 461
Watanabe, K., see Ishibashi, M. 183
Wetzel, D. L., see Pussayanawin, V. 243
Wigilius, B.
——, Borén, H., Carlberg, G. E., Grimvall, A., Lundgren, B. V. and Sävenhed, R.
Systematic approach to adsorption on XAD-2 resin for the concentration and analysis of trace organics in water below the μg/l level 169
Wilson, T. D.
High-performance liquid chromatographic determination of Tornalate® in solution dosage forms; a specificity study 409
Winzor, D. J., see Bergman, D. A. 67
Yamasaki, Y., see Kato, Y. 395
Yamashita, K., see Ishibashi, M. 183
Yang, Y.-B.
—— and Verzele, M.
High-speed and high-performance size-exclusion chromatography of proteins on a new hydrophilic polystyrene-based resin 383
Yeung, E. S., see Chan, K. C. 465
Yndestad, M., see Rogstad, A. 328
Ziegler, S. J.
——, Meier, B. and Sticher, O.
Rapid and sensitive determination of dehydroascorbic acid in addition to ascorbic acid by reversed-phase high-performance liquid chromatography using a post-column reduction system 419